

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE VETERINARY MEDICINAL PRODUCT**

Endofluke 100 mg/ml Oral Suspension

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

**Active substance:** per ml

Triclabendazole 100mg

**Excipients:**

Methyl Parahydroxybenzoate (E218)	2mg (preservative)
Propyl Parahydroxybenzoate (E216)	0.2mg (preservative)

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Oral Suspension.  
A white to off-white suspension.

### **4. CLINICAL PARTICULARS**

#### **4.1 Target species**

Cattle & Sheep

#### **4.2 Indications for use, specifying the target species**

For the treatment of adult, immature and early immature stages of liver fluke (*Fasciola hepatica*) susceptible to triclabendazole.

#### **4.3 Contraindications**

Do not use in animals known to be hypersensitive to the active substance .

#### **4.4 Special warnings for each target species**

Care should be taken to avoid the following practices, because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

- Too frequent and repeated use of anthelmintics from the same class, over an extended period of time.
- Under dosing, which may be due to underestimation of bodyweight, misadministration of the product or lack of calibration of the dosing device (if any).

Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of

the test(s) strongly suggest resistance to a particular anthelmintic, an anthelmintic belonging to another pharmacological class and having a different mode of action should be used.

Resistance to triclabendazole has been reported in *Fasciola hepatica* in cattle and sheep. Therefore, the use of this product should be based on local (regional/farm) epidemiological information about susceptibility of the *Fasciola hepatica* and recommendations on how to limit further selection for resistance to anthelmintics.

#### **4.5 Special precautions for use**

##### **i) Special precautions for use in animals**

Care should be taken when dosing animals to avoid causing injury to the mouth and pharynx.

##### **ii) Special Precautions to be taken by the person administering the product to animals**

When using, do not eat, drink or smoke.

Wash splashes from eyes and skin immediately.

Take off immediately any contaminated clothing.

Wash hands and exposed skin before meals and after use.

##### **iii) Other precautions**

The use of this product may have harmful effects on fish and aquatic invertebrates. Cattle and sheep must not have any access to the surface water such as streams, ponds or ditches within 7 days after treatment. When spreading manure from treated animals on arable lands, a safety distance of 10m to adjacent surface waters must be kept.

#### **4.6 Adverse reactions (frequency and seriousness)**

Occasionally, inflammation of the unpigmented skin, including the udder and the teats may occur after treatment in cattle exposed to intense sunshine.

#### **4.7 Use during pregnancy, lactation or lay**

The product is safe for use during pregnancy and lactation. However, the product is not permitted for use during lactation in animals producing milk for human consumption (see section 4.11).

#### **4.8 Interaction with other medicinal products and other forms of interaction**

None known

#### **4.9 Amounts to be administered and administration route**

For single oral administration only using properly calibrated dosing equipment. The product is suitable for most types of automatic drenching guns.

Shake the container before use.

Use unaltered from original container.

Clean drenching equipment before and after use.

**Dosage:**

Endofluke 100 mg/ml is given as an oral drench and is suitable for most types of automatic drenching guns.

The recommended dose rate is 12mg triclabendazole per kg bodyweight in cattle and 10mg triclabendazole per kg bodyweight in sheep.

**Practical Dosage Guide: Cattle: 6 ml per 50kg bodyweight**

Animal Weight	Dose of product
50kg	6ml
100kg	12ml
150kg	18ml
200kg	24ml
250kg	30ml
300kg	36ml
350kg	42ml
400kg	48ml
For each additional 50kg	6ml

**Sheep: 1 ml per 10kg bodyweight**

Animal Weight	Dose of product
10kg	1ml
20kg	2ml
30kg	3ml
40kg	4ml
50kg	5ml
60kg	6ml
For each additional 10kg	1ml

To ensure administration of a correct dose, bodyweight should be determined as accurately as possible; accuracy of the dosing device should be checked.

If animals are to be treated collectively rather than individually, they should be grouped according to their bodyweight and dosed accordingly, in order to avoid under- or overdosing.

The timing for re-treatment should be based on epidemiological risk patterns and should be customised for each individual farm.

To avoid the potential for the accumulation of residues following repeat administration of the product; animals should not be treated with a frequency of less than 10 weeks.

**4.10 Overdose**

A single oral dose of 150-200 mg triclabendazole/kg of live bodyweight may lead to side effects such as unsteady gait, dullness and reduced appetite. These side effects are slight and last 1 to 5 days. An antidote is not known.

#### **4.11 Withdrawal period(s)**

Cattle ( meat and offal): 56 days

Milk:

Cattle (milk): Milk for human consumption may only be taken from 48 hours after calving. Not intended for use within 45 days of calving. Should a cow calve earlier than 45 days after the last treatment, milk for human consumption may only be taken from 45 days + 48 hours (47 days) after the last treatment.

Sheep (meat & offal): 56 days

Sheep (milk): Not authorised for use in ewes producing milk for human consumption including during the dry period. Do not use within 1 year prior to the first lambing in ewes intended to produce milk for human consumption.

### **5. PHARMACOLOGICAL PROPERTIES**

Pharmacotherapeutic group: anthelmintics; benzimidazoles and related substances.  
ATCvet code: QP52ACO1

#### **5.1 Pharmacodynamic properties**

Triclabendazole differs from other benzimidazoles in that it is a narrow spectrum anthelmintic. The drug accumulates significantly in both immature and adult stages of *Fasciola hepatica* and stimulates the major routes of the parasite's energy generating system, as demonstrated by glucose derived acetate and propionate formation. However, under these conditions the parasite's motility decreased, indicating that the drug is not associated with inhibition of the energy generating pathways. Triclabendazole inhibits colchicine binding to microtubular proteins suggesting interference of the drug with microtubular structure and function. The drug strongly inhibits the release of proteolytic enzymes in immature and adult parasites, a process dependant on microtubular functions. The precise molecular mode of action of this fasciolicidal drug remains to be elucidated.

#### **5.2 Pharmacokinetic particulars**

50-75% of the orally administered dose of triclabendazole is absorbed from the gastrointestinal tract. Very rapidly, absorbed triclabendazole is almost completely oxidised to its sulfoxide and sulfone. In cattle triclabendazole sulfoxide reaches peak concentrations approximately 27 hours after administration of the product and the sulfone reaches peak concentrations 64 to 72 hours after administration. In sheep triclabendazole sulfoxide reaches peak concentrations approximately 20 hours after administration of the product and the sulfone reaches peak concentrations 30 to 32 hours after administration. Both metabolites bind strongly to plasma proteins, particularly albumin. Metabolites are excreted via the bile mainly as conjugates. More than 90%- 95% of the total dose of triclabendazole is excreted in the faeces, about 2% in the urine and less than 1% in the milk. The elimination is virtually complete by 10 days after administration.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Xanthan Gum  
Methyl Parahydroxybenzoate  
Propyl Parahydroxybenzoate  
Citric Acid Anhydrous  
Sodium Citrate  
Polysorbate 80  
Silica Colloidal  
Anhydrous Dimethicone Emulsion  
Water, purified

### **6.2 Incompatibilities**

None known.

### **6.3 Shelf life**

Shelf life of the veterinary medicinal product as packaged for sale: 2 years

### **6.4 Special precautions for storage**

Protect from frost.

### **6.5 Nature and composition of immediate packaging**

1 litre, 2.5 litre, 5 litre, 6 litre (5 litre +1 litre) and 7.5 litre (5 litre+2.5 litre) and 15 litre (3x5 litre) pack sizes high-density polyethylene flat bottom backpack containers containing a white to off-white smooth suspension. The closures used for all pack sizes are 38 mm propylene standard caps with an induction heat seal liner. A 38 mm polypropylene spouted cap is also provided with each pack size for dispensing purposes.

### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

The product may have toxic effects on fish and aquatic invertebrates. Any unused product or waste material must not enter surface water and should be disposed of in accordance with national requirements.

## **7. MARKETING AUTHORISATION HOLDER**

Bimeda Animal Health Limited  
2 / 3 / 4 Airton Close  
Tallaght  
Dublin 24  
Ireland

**8. MARKETING AUTHORISATION NUMBER**

Vm 50146/4018

**9. DATE OF FIRST AUTHORISATION**

20 February 2003

**10. DATE OF REVISION OF THE TEXT**

October 2018

**PROHIBITION OF SALE, SUPPLY AND/OR USE**

Not applicable.

Approved: 19 October 2018

A handwritten signature in black ink, appearing to read 'D. Austin', with a horizontal line extending from the end of the signature.